your peak flow diary



Recording your peak flow scores every day helps you to stay in control of your asthma and quickly spot if anything changes.



asthma with you every breath of the way



What is peak flow?

Peak flow is a measurement of how quickly you can blow air out of your lungs. You do this by blowing very hard into a plastic tube called a peak flow meter. Your GP or asthma nurse can show you how best to do this. You can buy a peak flow meter from most pharmacies or you can get one on prescription.

Your peak flow score

Peak flow scores will vary depending on your age, your height and whether you're a man or a woman. But it's the pattern that your scores make that's important, rather than one score on its own.

It's important to know your best peak flow. This will help you keep on top of your asthma. It's also helpful for when you visit your GP or asthma nurse as it will let you both assess your asthma.

Why is measuring peak flow useful?

Using the peak

flow diary every

day took less time

than brushing my

to ensure that my

disrupt my life.

teeth, and helps


asthma doesn't

- ✓ To tell what's going on in your lungs rather than guessing by how you feel
- Find out if your medicine is working or if you need to change the amount you're taking
- Keep a record of how well you've been and show this to your GP or asthma nurse
- Act quickly if your peak flow tells you it's an asthma emergency

How to take your peak flow

Blow into your peak flow meter first thing in the morning and in the evening before you take your asthma medicine.

When you blow into your meter do it three times and record the **highest** score with a cross on the chart inside.




How to record your peak flow

There are three sections to fill in:

- If you experience one of these warning symptoms, mark a cross in the box for that day. If you have marked at least one box a day for three consecutive days then see your GP or asthma nurse.
- Twice a day, record your highest peak flow score (after blowing into your meter three times) with a cross on the chart.
- Use this section to write down anything unusual or different that may have been the reason for a lower than normal peak flow in a given week, eg "I was stressed", "I exercised on Tuesday", "I came into contact with a pet".

Now you can start recording your peak flow overleaf



You can download more graph paper from www.asthma.org.uk/yourasthma

WHAT TO DO IF YOU HAVE AN ASTHMA ATTACK

An emergency is when any of the following happen:

- 1 Your reliever (blue) inhaler doesn't help
- 2 Your symptoms get worse
- **3** You are too breathless to speak
- 4 Your peak flow is only half what you normally blow (insert number here)

What to do in an asthma attack

- 1 Take two puffs of my reliever inhaler (one puff at a time)
- $\mathbf{2}~$ Sit up and try to take slow, steady breaths
- If I don't start to feel better, take two puffs of my reliever inhaler (one puff at a time) every two minutes. I can take up to ten puffs
- If I don't feel better I should call 999 straight away. If an ambulance doesn't arrive within ten minutes, and I'm still not feeling better, then I should repeat Step 3
- Even if I feel better after this I should see my GP or asthma nurse for advice the same day
- 6 If I have rescue prednisolone tablets, take 40mg (8 x 5mg) altogether

Please note this asthma attack information is not designed for people who use the Symbicort SMART regime. If you use Symbicort SMART please speak to your GP or asthma nurse about this.

Asthma UK has a range of resources to help with your asthma and a team of specialist asthma nurses if you need further advice.

Asthma UK Adviceline 0800 121 62 44

info@asthma.org.uk

Or visit our website www.asthma.org.uk



with you every breath of the way



@ 2013 Asthma UK registered charity number in England and Wales 802364 and in Scotland SC039322 HP2260513 First published 2005; Last updated 2011; Last reviewed 2013; Next review 2016